

Last updated: 1/10/22

Out-patient Treatment & Prophylaxis for COVID 19

Prophylaxis

Treatment Infusions

Treatment Oral

	EVUSHELD as Pre Exposure Prophylaxis <i>IM mAb</i>	MONOCLONAL Abs as Post Exposure Prophylaxis* <i>Subq injection</i>	MONOCLONAL Abs <i>RegenCOV / Sotrovimab / Bam/Ete</i> <i>Single infusion</i>	REMDESEVIR <i>Once daily Infusion x3</i>	MOLNUPIRAVIR	PAXLOVID <i>Nirmatrelvir + Ritonovir</i>
Intended Use	Passive introduction of neutralizing mAbs before illness or exposure	Passive introduction of neutralizing mAbs after known exposure in high risk individuals/environments (communal living/SNF/ etc.)	Passive introduction of neutralizing mAbs to reduce progression of disease and hospitalizations	Antiviral intended to reduce viral load early in viremia for those with mild/moderate symptoms	Antiviral intended to reduce viral load early in viremia to reduce risk of hospitalization	Boosted Protease Inhibitor intended to reduce viral load early in viremia to reduce risk of hospitalization
Who is eligible?	1. Severe allergy to COVID 19 vaccination OR 2. Profound immunocompromised innate or medically induced and unlikely mount appropriate response to vaccination	COVID negative individuals with known exposure in environment at high risk for an outbreak	Minimum criteria outlined in respective playbooks found on COVID 19 toolkit *eligibility may vary based on availability of product	Unvaccinated COVID positive individuals with clinical risk factors for progression to severe disease	COVID positive individuals with mild/moderate symptoms	COVID positive individuals with mild/moderate symptoms
When to prescribe?	When patient is well, requires negative COVID test *can NOT have active COVID*	Known positive in a high risk setting with outbreak potential (rehab, SNF, congregate living) OR Exposed COVID negative close contact with risk of severe disease	Early in symptomatic course for patient with documented positive SARS CoV 2 test and meets above criteria	Early in symptomatic course for patient with documented positive SARS CoV 2 test and meets above criteria	Early in symptomatic course, within 5 days of symptom onset and within 7 days of testing positive	Early in symptomatic course, within 5 days of symptom onset and within 7 days of testing positive
Misc.		This prophylaxis is variant specific and may not always be appropriate	mAb infusions are variant specific and will vary based on epidemiologic prevalence and availability		Through allocation to limited outpatient pharmacies, ordered through standard ordering processes. Several restrictions in special populations see pages 8-11: Special Considerations	Through allocation to limited outpatient pharmacies, ordered through standard ordering processes. Risk for drug drug interaction, see page 6: Contraindications

Last updated: 1/10/22

Ordering Work Flow

Prophylaxis

Treatment

Infusions

Treatment

Oral

EVUSHELD as Pre Exposure Prophylaxis
IM mAb

MONOCLONAL Abs as Post Exposure Prophylaxis*
Subq injection

MONOCLONAL Abs

RegenCOV / Sotrovimab / Bam/Ete
Single infusion

REMDESEVIR

Once daily Infusion x3

MOLNUPIROVIR

PAXLOVID

Nirmatrelvir + Ritonovir

Ordered Through

SLH Hospital pharmacies
Per standard immunization process

SLH Hospital pharmacy
Per relevant Playbook process

SLH Hospital pharmacies
Per relevant Playbook process

SLH Hospital pharmacies
Per relevant Playbook process

Limited community pharmacies

Limited community pharmacies

How to Order

1. Refer to Evusheld Playbook on Toolkit
2. Ordered by office for product to be handled and stored in appropriate clinic
3. For patient administration, drug available in standard ECW ordering process (does NOT require fax to designated pharmacy)

1. Refer to relevant Playbook for specific instructions
2. Orderable through ECW
3. Alternative paper order process available

1. Standard ordering through ECW
2. Receiving pharmacy **MUST** be one of the following:

- a) Kinney Drugs 17 Clinton St, Gouverneur
- b) The Medicine Place/Kimros 511 State St, Ogdensburg
- c) Adirondak Pharmacy 4057 State Highway 3, Star Lake

1. Standard ordering through ECW
2. Receiving pharmacy **MUST** be one of the following:

- a) Kinney Drugs 17 Clinton St, Gouverneur
- b) The Medicine Place/Kimros 511 State St, Ogdensburg
- c) Adirondak Pharmacy 4057 State Highway 3, Star Lake

Administration/ Handling

Given in outpatient clinic settings per standard IM injection practices and protocols

Administered at each SLH hospital in areas appropriate for infusion

Administered at each SLH hospital in areas appropriate for infusion

To be administered at each of the SLH hospitals in areas appropriate for infusion